

The Law Project for Psychiatric Rights (PsychRights®)

| Drug | Indication (diagnosis) | FDA Approval | DRUGDEX Support for Off-Label Use | DRUGDEX Recommendation Level |
|-----------------------|---|--|-----------------------------------|---|
| 2.49 | (Lingstone) | . 2777. pp . 0 va. | | |
| Key: | White Background: Medically Accepted Indication | | | |
| | Yellow Backgroun: Medically Accepted Indication Status Not Ascertained Orange Background: Pediatric Indication cited, but not supported by DRUGDEX | | | |
| | | | | X |
| | Red Background: No Pediatric FDA Ap | oproval or DRUGDEX c | itation | |
| Abilify (Aripiprazole | e) - Antipsychotic | | | |
| | Bipolar I Disorder - Adjunctive therapy with | | | |
| | lithium or valproate for Acute Manic or Mixed | Vac (for 40 um old and um) | | |
| | Episodes Bipolar I Disorder, monotherapy, Manic or Mixed | Yes (for 10 yrs old and up) Yes (for 10-17 years old re | | |
| | Episodes | acute therapy) | | |
| | Schizophrenia | Yes (for 13-17 years old) | | |
| Adderall (amphetar | nine/dextroamphetamine) - Central Nervous System Agen | • | | |
| (| | Yes (for 3 years old and up | | |
| | | re: [immediate-release] and | | |
| | | 6 years old and up re: | | |
| | Attention Deficit Hyperactivity Disorder (ADHD) | [extended-release] drug | | |
| | | Yes (for 6 years old and up re: [immediate release] | | |
| | Narcolepsy | drug) | | |
| Anafranil (clomip | ramine) - Antidepressant; Antidepressant, Tricyclic; Centr | ral Nervous System Agent | | |
| | | , , | | |
| | Depression | No | | Class IIb |
| [~· | Obsessive-Compulsive Disorder | Yes (for 10 years and up) | | |
| Clorazil (clozapine) | - Antipsychotic; Dibenzodiazepine | | | |
| | Bipolar I Disorder | No | | Class IIb |
| | | | | |
| | | | | cited, with no |
| | Cabinarda Tarata and Basistant | NI- | | recommendation |
| Caracarta | Schizophrenia, Treatment Resistant | No | | · · · · · · · · · · · · · · · · · · · |
| Concerta (methylph | Schizophrenia, Treatment Resistant nenidate) - Amphetamine Related; Central Nervous System | n Agent; CNS Stimulant | | recommendation |
| Concerta (methylph | nenidate) - Amphetamine Related; Central Nervous System | Agent; CNS Stimulant Yes (for 6 years old to 12 | | recommendation |
| Concerta (methylph | | n Agent; CNS Stimulant | | recommendation |
| Concerta (methylph | Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) | Agent; CNS Stimulant Yes (for 6 years old to 12 years old) Yes (for 6 years old and up) re ConcertaR | | recommendation |
| Concerta (methylph | Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Autistic Disorder | Agent; CNS Stimulant Yes (for 6 years old to 12 years old) Yes (for 6 years old and up) | | recommendation |
| Concerta (methylph | Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Autistic Disorder Impaired Cognition - inding related to | Agent; CNS Stimulant Yes (for 6 years old to 12 years old) Yes (for 6 years old and up) re ConcertaR No | | recommendation level |
| Concerta (methylpl | Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Autistic Disorder Impaired Cognition - inding related to coordination/ in coordination | Agent; CNS Stimulant Yes (for 6 years old to 12 years old) Yes (for 6 years old and up) Yes (for 6 years old and up) The Concertar No No | | recommendation level Class IIb Class IIb |
| Concerta (methylpl | Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Autistic Disorder Impaired Cognition - inding related to coordination/ in coordination Schizophrenia | n Agent; CNS Stimulant Yes (for 6 years old to 12 years old) Yes (for 6 years old and up) Te ConcertaR No No No | | recommendation level Class IIb Class IIII |
| | Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Autistic Disorder Impaired Cognition - inding related to coordination/ in coordination Schizophrenia Traumatic Brain Injury | n Agent; CNS Stimulant Yes (for 6 years old to 12 years old) Yes (for 6 years old and up) re ConcertaR No No No | | recommendation level Class IIb Class IIII Class IIII |
| Cymbalta (duloxeti | Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Autistic Disorder Impaired Cognition - inding related to coordination/ in coordination Schizophrenia Traumatic Brain Injury ine) - Antidepressant; Central Nervous System Agent; Neurone | n Agent; CNS Stimulant Yes (for 6 years old to 12 years old) Yes (for 6 years old and up) Yes (for 6 years old and up) Te ConcertaR No No No No No No Tropathic Pain Agent; Serotonin/I | Norepinephrine Reuptake | recommendation level Class IIb Class IIII Class IIII |
| Cymbalta (duloxeti | Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Autistic Disorder Impaired Cognition - inding related to coordination/ in coordination Schizophrenia Traumatic Brain Injury Ine) - Antidepressant; Central Nervous System Agent; Neuacid) – Anticonvulsant; Antimigraine; Valproic Acid (class | n Agent; CNS Stimulant Yes (for 6 years old to 12 years old) Yes (for 6 years old and up) Yes (for 6 years old and up) Te ConcertaR No No No No No No Tropathic Pain Agent; Serotonin/I | Norepinephrine Reuptake | recommendation level Class IIb Class IIII Class IIII |
| Cymbalta (duloxeti | Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Autistic Disorder Impaired Cognition - inding related to coordination/ in coordination Schizophrenia Traumatic Brain Injury ine) - Antidepressant; Central Nervous System Agent; Neurone | n Agent; CNS Stimulant Yes (for 6 years old to 12 years old) Yes (for 6 years old and up) Yes (for 6 years old and up) Te ConcertaR No No No No No No Tropathic Pain Agent; Serotonin/I | Norepinephrine Reuptake | recommendation level Class IIb Class IIII Class IIII |
| Cymbalta (duloxeti | Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Autistic Disorder Impaired Cognition - inding related to coordination/ in coordination Schizophrenia Traumatic Brain Injury Ine) - Antidepressant; Central Nervous System Agent; Neu acid) – Anticonvulsant; Antimigraine; Valproic Acid (class Absence Seizure, Simple and Complex and/or | n Agent; CNS Stimulant Yes (for 6 years old to 12 years old) Yes (for 6 years old and up) re ConcertaR No No No No No ropathic Pain Agent; Serotonin/ | Norepinephrine Reuptake | recommendation level Class IIb Class IIII Class IIII |
| Cymbalta (duloxeti | Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Attention Deficit Hyperactivity Disorder (ADHD) Autistic Disorder Impaired Cognition - inding related to coordination/ in coordination Schizophrenia Traumatic Brain Injury ine) - Antidepressant; Central Nervous System Agent; Neu acid) – Anticonvulsant; Antimigraine; Valproic Acid (clas Absence Seizure, Simple and Complex and/or Complex Partial Epileptic Seizure | n Agent; CNS Stimulant Yes (for 6 years old to 12 years old) years old) res (for 6 years old and up) re ConcertaR No No No No No ropathic Pain Agent; Serotonin/liss) Yes (10 years and older) | Norepinephrine Reuptake | recommendation level Class IIb Class III Class IIII Class IIIb Inhibitor |

1

The Law Project for Psychiatric Rights (PsychRights®)

| Drug | Indication (diagnosis) | FDA Approval | DRUGDEX Support | DRUGDEX Recommendation Level |
|--|---|---|--|------------------------------------|
| Dexedrine (dextroamphetamine) - Amphetamine (class); CNS Stimulant | | | | |
| | Attention Deficit Hyperactivity Disorder (ADHD) | Yes (for 3 years to 16 years old (immediate-release) and age 6 years to 16 years old (sustained-release)) | | |
| | Narcolepsy | | | |
| Desyrel (trazadone) - | Antidepressant; Triazolopyridine | Yes (for 6 years old and up) | | |
| Effexor (venlafaxine) – | Antidepressant; Antidepressant, Bicyclic; Phenethylan | nine (class); Serotonin/ Norepine | phrine Reuptake Inhibitor | ſ |
| | Attention Deficit Hyperactivity Disorder (ADHD) | No | | Class IIb |
| | Generalized Anxiety Disorder | No | | Class IIb |
| | Major Depressive Disorder | No | | Class IIb |
| | Severe Major Depression with Psychotic Features | "See Drug Consult Reference DEPRESSION - DRUG THE | | Chiss Ho |
| | Social Phobia | No | | Class IIb |
| Focalin (dexmethylphen | nidate) - Amphetamine Related; CNS Stimulant | | | |
| | Attention Deficit Hyperactivity Disorder (ADHD) | Yes (for 6 years and older) | | |
| Geodon (ziprasidone) - | Antipsychotic; Benzisothiazoyl | | | |
| <u>Haldol</u> (haloperidol) - | Antipsychotic; Butyrophenone; Dopamine Antagonis | | | |
| | Agitation | No | | Class IIb |
| | Hyperactive Behavior, (Short-term treatment) after failure to respond to non-antipsychotic medication and psychotherapy | Yes (for 3 years old and up) | (decanoate) is FDA approved for any pediatric use. DRUGDEX says safety and efficacy not established. | |
| | Problematic Behavior in Children (Severe), With failure to respond non-antipsychotic medication or psychotherapy | Yes (for 3 years old and up) | | |
| | Psychotic Disorder | Yes (for 3 years old and up but ORAL formulations only) | | |
| | Schizophrenia | Yes (for 3 years old and up but ORAL formulations only) | | |
| <u>Invega</u> (paliperidone) - | · Antipsychotic; Benzisoxazole | | | |
| <u>Lamictal</u> (lamotrigine) | - Anticonvulsant; Phenyltriazine | | | |
| | Bipolar Disorder, Depressed Phase | No | | Class IIb |
| | Epilepsy, Refractory | No | Class IIa | |
| Lexapro (escitalopram)- | Antianxiety, Antidepressant, Serotonin Reuptake Inhib | | | |
| | Major Depressive Disorder | Yes (for 12 years old and up) | | |
| <u>Luvox</u> (fluvoxamine) - | Antidepressant; Central Nervous System Agent; Serote | onin Reuptake Inhibitor | | |
| | Asperger's Disorder | No | | Class IIb |
| | Obsessive-Compulsive Disorder | Yes (for 8 years old and up and immediate release formula only) | | |
| | Severe Major Depression with Psychotic Features | "See Drug Consult Reference DEPRESSION - DRUG THE | | |

The Law Project for Psychiatric Rights (PsychRights®)

| Drug | Indication (diagnosis) | FDA Approval | DRUGDEX Support for Off-Label Use | DRUGDEX Recommendation Level |
|------------------------|--|--|-----------------------------------|------------------------------------|
| Mellaril (thioridazin | e) - Antipsychotic; Phenothiazine; Piperidine | | | |
| | Behavioral Syndrome | No | | Class IIII |
| | Schizophrenia, Refractory | Yes | | |
| Orap (pimozide) - A | antipsychotic; Diphenylbutylpiperidine; Dopamine Antago | onist | | |
| Paxil (paroxetine) - A | Antidepressant; Central Nervous System Agent; Serotonia | n Reuptake Inhibitor | | |
| Prozac (fluoxetine) - | Antidepressant; Central Nervous System Agent; Seroton | in Reuptake Inhibitor | | |
| | Anxiety Disorder of Childhood | No | | Class IIb |
| | Major Depressive Disorder | Yes (for 8 years old and up) | | |
| | Obsessive-Compulsive Disorder | Yes (for 7 years old and up | | |
| | Severe Major Depression with Psychotic Features | "See Drug Consult Reference DEPRESSION - DRUG THE | | |
| Ritalin (methylpheni | date) - Amphetamine Related; Central Nervous System A | * | | |
| | Attention Deficit Hyperactivity Disorder (ADHD) | Yes (for 6 years to 12 years old)(exteded release) | | |
| | Attention Deficit Hyperactivity Disorder (ADHD) | Yes (for 6 years old and up)(immediate release) | | |
| | Narcolepsy | Yes (for 6 years and up, and Ritalin(R) -SR only) | | |
| | Schizophrenia | No | | Class IIII |
| Risnerdal (risperide | Traumatic Brain Injury one) - Antipsychotic; Benzisoxazole | No | | Class IIb |
| (Hisperial | Sile) - Antipsychotic, Benzisokuzote | | | |
| | Autistic Disorder – Irritability | Yes (for 5 years old and up) | | |
| | Bipolar I Disorder | Yes (for 10 years old and up) | | |
| | Schizophrenia | Yes (for 13 years old and up, ORALLY) | | |
| Seroquel (QUETIA | PINE) - Antipsychotic; Dibenzothiazepine | | | |
| | Schizophrenia | No | | Class IIb |
| Sinequan (doxepin) | - Antianxiety Antidepressant; Antidepressant, Tricyclic; | | | |
| | Alcoholism - Anxiety – Depression | Yes (for 12 years old and up) | | |
| | Anxiety – Depression | Yes (for 12 years old and up) | | |
| | Anxiety - Depression - Psychoneurotic personality disorder | Yes (for 12 years old and up) | | |

The Law Project for Psychiatric Rights (PsychRights®)

| Drug | Indication (diagnosis) | FDA Approval | DRUGDEX Support for Off-Label Use | DRUGDEX Recommendation Level |
|----------------------------|--|---|-----------------------------------|------------------------------------|
| Strattera (atomoxetine | | | | |
| | | | | |
| | Attention Deficit Hyperactivity Disorder (ADHD) | Yes (for 6 years old and up) | | |
| | Attention Deficit Hyperactivity Disorder (ADHD) - | | | |
| G 1 | Social phobia | No | | Class IIb |
| | ydrochloride/olanzapine) - Antidepressant; Antipsycho | | | |
| Tegretol (carbamazep | ine) - Anticonvulsant; Antimanic; Dibenzazepine Carb | oxamide; Neuropathic Pain Ager | nt | |
| | Epilepsy, Partial, Generalized, and Mixed types | Yes | | |
| | Migraine; Prophylaxis | | | Class IIb |
| <u> </u> | Neuropathy, General | | | Class IIb |
| Tofranil (imipramine) | - Antidepressant; Antidepressant, Tricyclic; Urinary F | Enuresis Agent | | |
| | Attention Deficit Hyperactivity Disorder (ADHD), | | | |
| | Predominantly Inattentive Type | No | | Class IIb |
| | Depression | No | | Class IIb |
| | | | | |
| | Nocturnal enuresis | Yes (for 6 years old and up) | | |
| | Separation Anxiety Disorder of Childhood | No | | Class IIII |
| Trilontal | Schizophrenia, Adjunct | No | | Class IIII |
| 1 rueptai (oxcarbazepi | ne) - Anticonvulsant; Dibenzazepine Carboxamide | | | |
| | Partial Seizure, monotherapy | Yes (for 4 years old and up) | | |
| Ī | Partial seizure; Adjunct | Yes (for 2 years old and up) | | |
| Vyvanse (lisdexamfeta | mine) - Amphetamine (class); CNS Stimulant | | | |
| | Attention Deficit Hyperactivity Disorder (ADHD) | Yes (for 6 years old to 12 years) | | |
| Zoloft (sertraline) - Anti | depressant; Central Nervous System Agent; Serotonin | Reuptake Inhibitor | | |
| | Obsessive-Compulsive Disorder | Yes (6 years old and up) | | |
| | Anorexia nervosa | No | | Class IIII |
| | Generalized Anxiety Disorder | No | | Class IIb |
| | Major Depressive Disorder | No | | Class IIb |
| | Severe Major Depression with Psychotic Features | "See Drug Consult Referenc DEPRESSION - DRUG THE | | |
| Zyprexa (olanzapine) - | Antipsychotic; Thienobenzodiazepine | | | |
| | Bipolar I Disorder, Acute Mixed or Manic Episodes | No | Class IIa | |
| | Pervasive Developmental Disorder | No | | Class IIb |
| | Schizophrenia | No | | Class IIb |
| | Schizophrenia, Refractory | No | | Class IIb |
| | Severe Major Depression with Psychotic Features | "See Drug Consult Referenc DEPRESSION - DRUG THE | | |

DRUGDEX® Consults

RECOMMENDATION, EVIDENCE AND EFFICACY RATINGS

<u>RESPONSE</u>
The Thomson Efficacy, Strength of Evidence and Strength of Recommendation definitions are outlined below:

| Table 1. Strength Of Recommendation | | | |
|-------------------------------------|-----------------------|--|--|
| Class I | Recommended | The given test or treatment has been proven to be useful, and should be performed or administered. | |
| Class IIa | | The given test, or treatment is generally considered to be useful, and is indicated in most cases. | |
| Class IIb | 1 | The given test, or treatment may be useful, and is indicated in some, but not most, cases. | |
| Class III | Not Recommended | The given test, or treatment is not useful, and should be avoided. | |
| Class Indeterminant | Evidence Inconclusive | | |

| Table 2. S | Table 2. Strength Of Evidence | | | |
|----------------|--|--|--|--|
| | Category A evidence is based on data derived from: Meta-analyses of randomized controlled trials with homogeneity with regard to the directions and degrees of results between individual studies. Multiple, well-done randomized clinical trials involving large numbers of patients. | | | |
| В | Category B evidence is based on data derived from: Meta-analyses of randomized controlled trials with conflicting conclusions with regard to the directions and degrees of results between individual studies. Randomized controlled trials that involved small numbers of patients or had significant methodological flaws (e.g., bias, drop-out rate, flawed analysis, etc.). Nonrandomized studies (e.g., cohort studies, case-control studies, observational studies). | | | |
| Category C | Category C evidence is based on data derived from: Expert opinion or consensus, case reports or case series. | | | |
| No Evidence | | | | |

| Table 3 | Table 3. Efficacy | | | |
|--------------|-----------------------------|--|--|--|
| Class I | Effective | Evidence and/or expert opinion suggests that a given drug treatment for a specific indication is effective | | |
| Class Ila | | Evidence and/or expert opinion is conflicting as to whether a given drug treatment for a specific indication is effective, but the weight of evidence and/or expert opinion favors efficacy. | | |
| Class Ilb | Evidence is Inconclusive | Evidence and/or expert opinion is conflicting as to whether a given drug treatment for a specific indication is effective, but the weight of evidence and/or expert opinion argues against efficacy. | | |
| Class III | Ineffective | Evidence and/or expert opinion suggests that a given drug treatment for a specific indication is ineffective. | | |

© 1974- 2008 Thomson Healthcare. All rights reserved.